



Diamond Quantum Nanoemitters 150113

Jeronimo Maze
PONTIFICIA UNIVERSIDAD CATOLICA DE CHILE

05/05/2017
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/ IOS
Arlington, Virginia 22203
Air Force Materiel Command

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Executive Services, Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</p>						
1. REPORT DATE (DD-MM-YYYY) 31-05-2017		2. REPORT TYPE Final		3. DATES COVERED (From - To) 01 Mar 2015 to 31 Aug 2016		
4. TITLE AND SUBTITLE Cross Discipline Research on Hyperbolic Optical Systems for Control of Quantum Nanoemitters				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER FA9550-15-1-0113		
				5c. PROGRAM ELEMENT NUMBER 61102F		
6. AUTHOR(S) Jeronimo Maze				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) PONTIFICIA UNIVERSIDAD CATOLICA DE CHILE AV. LIBERTADOR BERNARDO OHIGG SANTIAGO, 8331150 CL				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/SOARD U.S. Embassy Santiago Av. Andres Bello 2800 Santiago, Chile				10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/AFOSR IOS		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-AFOSR-CL-TR-2017-0006		
12. DISTRIBUTION/AVAILABILITY STATEMENT A DISTRIBUTION UNLIMITED: PB Public Release						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT <p>As final results of this project there are: (1) the investigation of the effect of phonons on the optical properties of solid state emitters. A microscopic model was developed to calculate the phononic spectral density function responsible for the optical line shape of colour centres in diamond and alike impurities in large-bandgap materials (PHYS REV B 94, 134305 (2016)); (2) the exploration of the coupling between strain and electronic charge in mechanical resonators using a single quantum emitter in diamond. The work was a collaboration with the Department of Physics at University of California, Santa Barbara. The work was published in PHYS. REV. APPL. 6, 0340055 (2016). (3) the investigation of several noise sources, magnetic and electric, on the relaxation time of electronic spins associate to colour centers in diamond. It was found that the relative sensitivity to magnetic and electric noise can be adjusted by external magnetic fields (PHYS REV B 93, 024305 (2016); (4) the use of electronic spins as sensors of the local nuclear spin polarisation in diamond. Novel spectroscopical techniques were developed in order to obtain useful information about the nuclear bath (PHYS REV B 92, 241117R (2015).</p>						
15. SUBJECT TERMS nanoemitters, SOARD						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON POKINES, BRETT	
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) (703) 835-2309	

SCIENTIFIC REPORT

Federal grant Identifying number	FA9550-15-1-0113
Recipient Organization	PONTIFICIA UNIVERSIDAD CATOLICA DE CHILE, AVENIDA VICUNA MACKENNA 4860, SANTIAGO 7820436, CHILE
Recipient Principal Investigator	Dr. Jeronimo Maze, Physics Faculty, Pontificia Universidad Catolica, 011.56.2.2354.4486, jmaze@uc.cl
DUNS	
EIN	
Period	JAN, 01, 2015 - JUN, 30, 2016
Reporting Period End Date	SEP, 30, 2016
Govt Program Manager	Dr. Brett Pokines, AFOSR/IO, 011.56.2.330.3184, brett.pokines@afosr.af.mil

1. Introduction

This 18-moth AFOSR project allowed us to explore several aspects of color centers in solids such as the engineering of the optical properties of single emitters in large band gap materials and the understanding of the effect of external perturbations such as electronic and magnetic noise on the relaxation process of electronic spins associated to impurities. Such understanding has allowed us to take advantage of the great control that is possible to achieve of the internal degrees of freedom and use them to implement novel sensors of small perturbations such as the magnetic field produced by nuclear spins and the strain caused by mechanical resonators that are made out of the same material that host these single emitters.

In what follows we list the main results and undergoing research.

2. Results

2.1 Microscopic modeling of the effect of phonons on the optical properties of solid-state emitters

Published in	PHYSICAL REVIEW B 94, 134305 (2016)
Authors	Ariel Norambuena [1,2], Sebastian A. Reyes [1,2], Jose Mejia-Lopez [1,2], Adam Gali, [3,4] and Jeronimo R. Maze[1,2]
Affiliations	[1] Faculty of Physics, Pontificia Universidad Cato'lica de Chile, Avda. Vicuña Mackenna 4860, Santiago, Chile [2] Center for Nanotechnology and Advanced Materials CIEN-UC, Pontificia Universidad Católica de Chile, Avda. Vicuña Mackenna 4860, Santiago, Chile [3] Department of Atomic Physics, Budapest University of Technology and Economics, Budafoki ut 8., H-1111 Budapest, Hungary.

	[4] Institute for Solid State Physics, Wigner Research Centre for Physics, Hungarian Academy of Sciences, P. O. Box 49, H-1525, Budapest, Hungary.
--	--

Summary

Understanding the effect of vibrations in optically active nanosystems is crucial for successfully implementing applications in molecular-based electro-optical devices, quantum information communications, single photon sources, and fluorescent markers for biological measurements. Here, we present a first-principles microscopic description of the role of phonons on the isotopic shift presented in the optical emission spectrum associated to the negatively charged silicon-vacancy color center in diamond. We use the spin-boson model and estimate the electron-phonon interactions using a symmetrized molecular description of the electronic states and a force-constant model to describe molecular vibrations. Group theoretical arguments and dynamical symmetry breaking are presented in order to explain the optical properties of the zero-phonon line and the isotopic shift of the phonon sideband.

See Appendix 1 for more information.

2.2 Strain Coupling of a Mechanical Resonator to a Single Quantum Emitter in Diamond

Published in	PHYSICAL REVIEW APPLIED 6, 034005 (2016)
Authors	Kenneth W. Lee [1], Donghun Lee [1,2], Preeti Ovarthaiyapong [1], Joaquin Minguzzi [3,] Jero R. Maze [3] and Ania C. Bleszynski Jayich[1]
Affiliations	[1] Department of Physics, University of California Santa Barbara, Santa Barbara, California 93106, USA [2] Department of Physics, Korea University, Seoul 02841, South Korea [3] Institute of Physics, Pontificia Universidad Católica de Chile, Santiago 7820436, Chile

Summary

The recent maturation of hybrid quantum devices has led to significant enhancements in the functionality of a wide variety of quantum systems. In particular, harnessing mechanical resonators for manipulation and control has expanded the use of two-level systems in quantum-information science and quantum sensing. Here, we report on a monolithic hybrid quantum device in which strain fields associated with resonant vibrations of a diamond cantilever dynamically control the optical transitions of a single nitrogen-vacancy (NV) defect center in diamond. We quantitatively characterize the strain coupling to the orbital states of the NV center and, with mechanical driving, we observe NV-strain couplings exceeding 10 GHz. Furthermore, we use this strain-mediated coupling to match the frequency and polarization dependence of the zero- phonon lines of two spatially separated and initially distinguishable NV centers. The experiments demonstrated here mark an important step toward engineering a quantum device capable of realizing and probing the dynamics of

nonclassical states of mechanical resonators, spin systems, and photons.

2.3 Competition between electric field and magnetic field noise in the decoherence of a single spin in diamond

Published in	PHYSICAL REVIEW B 93, 024305 (2016)
Authors	P. Jamonneau[1], M. Lesik[1], J. P. Tetienne[1], I. Alvizu[2], L. Mayer[1], A. Dreau[1], S. Kosen[1], J.-F. Roch[1], S. Pezzagna[3], J. Meijer[3], T. Teraji[4], Y. Kubo[5], P. Bertet[5], J. R. Maze[2] and V. Jacques[1,6]
Affiliations	[1] Laboratoire Aime Cotton, CNRS, Universite Paris-Sud, ENS Cachan, Universite Paris-Saclay, 91405 Orsay Cedex, France [2] Facultad de Fisica, Pontificia Universidad Catolica de Chile, Santiago 7820436, Chile [3] Department of Nuclear Solid State Physics, Institute for Experimental Physics II, Universitat Leipzig, Linnzstrasse 5, 04103 Leipzig, Germany [4] National Institute for Materials Science, 1-1 Namiki, Tsukuba, Ibaraki 305-0044, Japan [5] Quantronics group, SPEC, CEA, CNRS, Universite Paris-Saclay, CEA Saclay 91191 Gif-sur-Yvette, France [6] Laboratoire Charles Coulomb, Universite de Montpellier and CNRS, 34095 Montpellier, France

Summary

We analyze the impact of electric field and magnetic field fluctuations in the decoherence of the electronic spin associated with a single nitrogen-vacancy (NV) defect in diamond. To this end, we tune the amplitude of a magnetic field in order to engineer spin eigenstates protected either against magnetic noise or against electric noise. The competition between these noise sources is analyzed quantitatively by changing their relative strength through modifications of the host diamond material. This study provides significant insights into the decoherence of the NV electronic spin, which is valuable for quantum metrology and sensing applications.

2.4 Local probing of nuclear bath polarization with a single electronic spin

Published in	PHYSICAL REVIEW B 92, 241117(R) (2015)
Authors	P. London [1] R. Fischer [1], I. Alvizu, [2], J. R. Maze [2] and D. Gershoni [1]
Affiliations	[1] Department of Physics, Technion, Israel Institute of Technology, Haifa 3200003, Israel [2] Departamento de Fisica, Pontificia Universidad Catolica de Chile, Santiago 7820436, Chile

Summary

The effect of a polarized nuclear spin bath on the dynamical behavior of a single electronic spin is studied theoretically and experimentally. The polarization of a single nuclear spin modifies the spin-echo signal of its neighboring electronic spin. When the electronic spin is surrounded by a bath of polarized nuclei, the spin-echo signals manifest a characteristic frequency related only to the nuclear spins abundance and their collective polarization. This frequency is proposed as an indicator for the local nuclear bath polarization. We quantify the realistic experimental regimes at which the scheme is efficient. Our proposal has potential applications for quantum sensing schemes, and opens a route for a systematic study of polarized mesoscopic systems.

3. Undergoing research

3.1 Use of diamond color centers for fluorescent markers of Amyloid beta fibrils

We are currently exploring the use of color centers in diamond to mark amyloid beta compounds, which are believed to be a precursor of the Alzheimer disease. We have successfully functionalized the surface of 30-nm diameter nanodiamonds in order to penetrate biological barriers and to detect amyloid beta fibrils in the extracellular region.

4. Concluding remarks

The AFOSR grant allowed us to explore several aspects of color centers in large bandgap materials. We developed a microscopic model to explore the effect of phonons on the optical properties of solid-state emitters. We calculated the phononic spectral density function responsible for the optical line shape of colour centres in diamond and alike impurities in large-bandgap materials. It was also possible to explore the coupling between strain and electronic charge in mechanical resonators using a single quantum emitter in diamond. We investigated several noise sources, magnetic and electric, on the relaxation time of electronic spins associate to color centers in diamond. It was found that the relative sensitivity to magnetic and electric noise can be adjusted by external magnetic fields. Finally, we use the electronic spin associate to the nitrogen-vacancy center in diamond as a sensor of the local nuclear spin polarisation in diamond. Novel spectroscopical techniques were developed in order to obtain useful information about the nuclear bath.

Jeronimo Maze
Associate Professor
Institute of Physics, Faculty of Physics
Pontificia Universidad Catolica de Chile

Appendix

A.1 Microscopic modeling of the effect of phonons on the optical properties of solid-state emitters

Here we describe the findings of the results published in PHYSICAL REVIEW B 94, 134305 (2016).

We have presented a microscopic model for estimating the emission spectrum of the SiV⁻ using the Kubo formula and the spin-boson model. In addition we have considered effects to second order on the spectral density function via dynamical symmetry breaking. This spectral density function is estimated using a force-constant model for describing the vibrational modes and symmetrized electronic wave functions constructed using group theoretical arguments. This approach allows us to gain detailed insight on the microscopic origin and the role of symmetries on the emission spectra and the spectral density function, an approach which is crucially different from, but validates, phenomenological models presented in previous works. These results might be useful for understanding and engineering the optical properties of color centers in solids by extending the analysis to other deep and shallow centers coupled to phonons and subject to instabilities such as dynamic Jahn-Teller effects and external perturbations such as electric fields or strain.




Figure A1. Schematic representation of the potential energy diagram. The two parabolas represent the phononic potential of the ground e_{gx} and excited e_{ux} states of the SiV⁻ including vibrational levels. Structure of the SiV⁻ in diamond: six carbon atoms (dark gray) and the interstitial silicon atom (green) embedded in a diamond lattice (light gray). The molecular orbital representation of the electronic states e_{gx} and e_{ux} are represented by red (blue) for the positive (negative) sign of the electronic wave function.




Figure A2. Numerical emission spectra of the SiV^- in diamond. The blue and red curves represent the numerical emission spectrum obtained for $T = 4 \text{ K}$ and $T = 296 \text{ K}$, respectively. The ZPL at 736 nm and the prominent sharp feature of the phonon sideband at 766 nm are reproduced. The peak at 766 nm is associated with the a_{1u} quasilocal phonon mode.